

UR-1353
ER-40685-804

**COMMENT ON “EQUIVALENCE OF SEVERAL
CHERN-SIMONS MATTER MODELS”**

by

C.R. Hagen
Department of Physics and Astronomy
University of Rochester
Rochester, NY 14627

It has recently been claimed¹ that there is a complete equivalence between three different models each of which consists of a matter field minimally coupled to a Chern-Simons gauge field as well as to an external vector field C_μ . These three cases consist of systems of spins $\frac{1}{2}$, 1, and $\frac{3}{2}$ but each with a different coefficient for the Chern-Simons term. The object of this note is to point out that such a claim is totally untenable.

It is simplest to begin by disposing of the spin- $\frac{3}{2}$ model. As is well known, a result of Johnson and Sudarshan² requires the matrix multiplying the time derivative term in the Lagrangian to be singular in order to have a positive definite metric³. Since the matrices of ref. 1 are simply the non-singular spin- $\frac{3}{2}$ representation of angular momentum, one concludes that Eq. (3) of ref. 1 describes an indefinite metric theory albeit with twice as many degrees of freedom as the single spin component models of Eqs. (1) and (2). Alternatively stated, since no one has succeeded to date in consistently quantizing the minimal spin- $\frac{3}{2}$ field coupled to a gauge field, it is quite impossible to establish an equivalence to the relatively well understood theories of Eqs. (1) and (2).

Next one turns to the equivalence claimed between the spinor and vector theories. The authors of ref. 1 discuss the current correlation functions at the one loop level in support of their result. It is known that this quantity is finite for all momenta p in the spinor case but divergent for the vector field⁴. In ref. 1 this complication is handled by the remark that dimensional regularization⁵ can eliminate the divergence at $p = 0$. No additional momentum dependence is calculated and it is merely hypothesized that all higher order corrections do not affect the conclusion. In fact one can readily calculate the $O(p^2/M^2)$ corrections to Π_e and Π_o using results of ref. 4. The answer is that the magnitude of the (finite) $O(p^2/M^2)$ correction to Π_o for the vector model is exactly twice that of the spinor model, a result which directly contradicts the equivalence claim of ref. 1.

Since the Aharonov-Bohm scattering problems for spin- $\frac{1}{2}$ ⁶ and spin one⁷ have been extensively studied, it is also of interest to make use of the fact that the results in these two cases are totally dissimilar. They can be realized in the context of ref. 1 by taking $\alpha \rightarrow \infty$ and C_μ to be the vector potential of a single flux tube. In the spinor case the solution is singular but normalizable even when the Zeeman term is attractive while the vector model has totally different properties. Significantly, there exists no state of lowest

energy for the latter when the spin is oriented parallel to the flux tube.

In summary then it is clear that the equivalence claims made in ref. 1 are totally incompatible with well known results.

This work was supported in part by U.S. Department of Energy Grant No. DE-FG-02-91ER40685.

References

1. W. Chen and C. Itoi, Phys. Rev. Lett. **72**, 2527 (1994).
2. K. Johnson and E.C.G. Sudarshan, Ann. Phys. (N.Y.) **13**, 126 (1961).
3. It is straightforward to verify that the theorem applies also to $2 + 1$ space.
4. C.R. Hagen, P. Panigrahi, and S. Ramaswamy, Phys. Rev. Lett. **61**, 389 (1988).
5. This would seem to raise the question as to why dimensional regularization should be used in this calculation since it is not required for the path integral derivation.
6. C.R. Hagen, Phys. Rev. Lett. **64**, 503 (1990) and **64** 2347 (1990).
7. C.R. Hagen and S. Ramaswamy, Phys. Rev. **D42**, 3524 (1990).